


CDC Data Show Almost 25,000 Adverse Events, 221 Deaths Reported After Bivalent COVID Boosters



Data from the Vaccine Adverse Event Reporting System (VAERS) released Friday show [1,533,182](#) adverse events following COVID-19 vaccines were reported between Dec. 14, 2020, and March 10, 2023. This includes [285,196](#) reports of serious injuries and [34,725](#) deaths.

Of the 34,725 reported deaths, [21,733 cases](#) are attributed to Pfizer, [9,770](#) to Moderna, [2,964](#) to Johnson & Johnson, and 18 to Novavax. Of the [reported deaths](#), 9% occurred within 24 hours of vaccination, and 13% occurred within 48 hours.

VAERS is a voluntary reporting system co-managed by the U.S. Food and Drug Administration and Centers for Disease Control and Prevention (CDC) designed to detect vaccine safety signals.

Excluding “[foreign reports](#)” to VAERS, [943,912 adverse events](#), including [17,114 deaths](#) and [102,050 serious injuries](#), were reported in the U.S. between Dec. 14, 2020, and March 10, 2023.

[Foreign reports](#) are reports from foreign subsidiaries sent to U.S. vaccine manufacturers. Under FDA regulations, if a manufacturer is notified of a foreign case report describing an event that is both serious and does not appear on the product’s labeling, the manufacturer must submit the report to VAERS.

In the U.S., 672 million [COVID-19 vaccine doses](#) had been administered as of March 8, including 401 million doses of Pfizer, 252 million doses of Moderna, 19 million doses of Johnson & Johnson, and 82,000 doses of Novavax.

Bivalent Booster Data

As of March 15, [54.2 million people](#) received an experimental bivalent booster dose targeting the no-longer-existing Wuhan strain and obsolete BA.4/BA.5 omicron subvariants.

Since the rollout of bivalent boosters in September 2022 and March 10, there have been [24,641 adverse events](#) reported to VAERS, with 40% attributed to [Moderna’s booster](#) and 60% attributed to [Pfizer/BioNTech](#). The data included [221 deaths](#), [1,853 serious injuries](#), and [73 reports](#) of myocarditis and pericarditis (heart inflammation).

The CDC uses a [narrowed case definition](#) of myocarditis. To meet the case definition of myocarditis, people must have had “symptoms such as chest pain, shortness of breath and feelings of having a fast-beating, fluttering or pounding heart, and medical tests to support the diagnosis of myocarditis and rule out other causes.” This allows them to exclude cases of cardiac arrest, ischemic strokes, and deaths due to heart problems that occur before one has the chance to go to the

hospital, obtain a diagnosis, or “dies suddenly.”

The CDC website does not state what happens to these cases, but there is no indication they are tracked or included in the CDC’s myocarditis numbers.

Data for 6-month-olds to 5-year-olds

- [5,950 adverse events](#), including [253 cases rated as serious](#) and [13 reported deaths](#).
- [5 reports](#) of myocarditis and pericarditis.
- [35 reports](#) of blood clotting disorders.
- [59 reports](#) of seizures.

Data for 5- to 11-year-olds

- [17,093 adverse events](#), including [829 rated as serious](#) and [34 reported deaths](#).
- [48 reports](#) of myocarditis and pericarditis.
- [76 reports](#) of blood clotting disorders.
- [193 reports](#) of seizures.

Data for 12- to 17-year-olds

- [41,527 adverse events](#), including [4,646 rated as serious](#) and [139 reported deaths](#).
- [276 reports](#) of anaphylaxis among 12- to 17-year-olds where the reaction was life-threatening, required treatment, or resulted in death.
- [1,352 reports](#) of myocarditis and pericarditis, with [1,184 cases](#) attributed to Pfizer’s vaccine.
- [312 reports](#) of blood clotting disorders, with [285 cases](#) attributed to Pfizer.
- [31 cases](#) of postural orthostatic tachycardia syndrome (POTS) were attributed to Pfizer’s vaccine.

Data for all age groups to VAERS

- 16% of deaths were related to cardiac disorders.
- 53% of those who [died were male](#), and 41% [were female](#). The remaining death reports do not list the gender of the deceased.
- The [average age](#) of death was 72.
- As of Feb. 17, [9,035 pregnant women](#) reported adverse events related to COVID-19 vaccines, including [5,231 reports](#) of miscarriage or premature birth.
- Of the [17,139 cases of Bell's palsy](#) reported, 73% were attributed to Pfizer vaccinations, [22% to Moderna](#), and [5% to J&J](#).
- [3,258 reports](#) of Guillain-Barré syndrome.
- [10,400 reports](#) of anaphylaxis where the reaction was life-threatening, required treatment or resulted in death.
- [8,488 reports](#) of myocardial infarction and cardiac arrest.
- [46,131 reports](#) of blood-clotting disorders. Of those, [31,677 reports](#) were attributed to Pfizer, [10,436 reports](#) to Moderna, and [3,943 reports](#) to Johnson & Johnson.
- [25,487 cases](#) of myocarditis and pericarditis, with [19,349 cases](#) attributed to Pfizer, [5,636 cases](#) to Moderna, and [442](#) to Johnson & Johnson.
- [79 cases](#) of Creutzfeldt-Jakob disease, with [65 cases](#) attributed to Pfizer, [12](#) to Moderna, and [2](#) to J&J.
- [706 cases](#) of POTS, with [523 cases](#) attributed to Pfizer, [155 cases](#) to Moderna, and [27 cases](#) to Johnson & Johnson.

Although healthcare providers are required by law to report vaccine adverse events to VAERS, research shows very few do. It is essential that anyone who experiences an adverse event [report their own injury](#).